

Please Note: This information represents many hours of diligent research over many years and is protected by copyright. Please do not copy more than 200 words, or photocopy all or part to pass on to others (either freely or for gain) without the express permission of the author, Jane Thurnell-Read

We do not believe that the written information alone provides sufficient information for accurate kinesiology testing or dowsing. Please do not use this information in that way.

LYME PLUS TEST KIT (61 Vials)

Product Code 8094

Please Note:

Some of the species included have only ever been documented in animals, but because differentiation of the various species is time-consuming and not always carried out when people present with symptoms, these species have been included.

There is some indication of countries where a particular species has been found, but this does not mean that it is only present in the countries listed.

Cat Scratch Disease

Lymph nodes, especially those around the head, neck, and upper limbs, become swollen. Additionally, a person with CSD may experience fever, headache, fatigue, and a poor appetite.

Lyme Disease

Typical symptoms include fever, headache, fatigue, and a characteristic skin rash called erythema migrans. If left untreated, infection can spread to joints, the heart, and the nervous system.

Relapsing Fever

Relapsing or recurring episodes of fever, often accompanied by headache, muscle and joint aches and nausea.

Product Code	Name	Comments
LY 01	Anaplasma Phagocytophilum / Ehrlichia Phagocytophilum	Causes human granulocytic anaplasmosis. Symptoms may include fever, severe headache, muscle aches (myalgia), chills and shaking, similar to the symptoms of influenza. GI symptoms occur in less than half of patients and a skin rash is seen in less than 10% of patients. It is also characterized by thrombocytopenia, leukopenia, and elevated serum transaminase levels in the majority of infected patients.
LY 02	Babesia Bigemina	North and South America, Southern Europe, Africa, Asia and Australia
LY 03	Babesia Bovis	Infects cattle and occasionally humans. Eradicated from the United States by 1943, but is still present in Mexico and much of the world's tropics.
LY 04	Babesia Canis	
LY 05	Babesia Cati	
LY 06	Babesia Divergens	Has been found in Turkey, Spain, Canary Islands, Tunisia, Austria, France and Norway. Infections have a much higher fatality rate (42%) than with other strains and present with the most severe symptoms: haemoglobinuria followed by jaundice, a persistently high fever, chills and sweats. If left untreated, can develop into shock-like symptoms with pulmonary oedema and renal failure.
LY 07	Babesia Duncanii	Can infect humans.
LY 08	Babesia Felis	

© Jane Thurnell-Read 1992-2012

Life-Work Potential, Sea View House, Long Rock, Penzance, TR20 8JF, UK

Tel UK: 0844 412 4487; Tel from outside UK: +44 1726 819179

office@lifeworkpotential.com | www.lifeworkpotential.com/

Product Code	Name	Comments
LY 09	Babesia Gibsoni	
LY 10	Babesia Herpailuri	
LY 11	Babesia Jakimoni	
LY 12	Babesia Major	
LY 13	Babesia Microti / Theileria Microti	Common in US. For 25% of cases in adults and half of cases in children, the disease is asymptomatic or mild with flu-like symptoms. Symptoms are characterized by irregular fevers, chills, headaches, general lethargy, pain and malaise.
LY 14	Babesia Ovate	
LY 15	Babesia Pantherae	
LY 16	Bartonella alsatica	
LY 17	Bartonella Arupensis	
LY 18	Bartonella Bacilliformis	Causes Carrion's disease (Oroya fever, Verruga peruana).
LY 19	Bartonella Berkhoffii	Becoming more important particularly for immuno-compromised individuals.
LY 20	Bartonella Birtlesii	
LY 21	Bartonella Bovis	
LY 22	Bartonella Capreoli	
LY 23	Bartonella Clarridgeiae	Found in domestic cats and can give humans Cat Scratch Disease
LY 24	Bartonella Doshiae	May cause Cat Scratch Disease.
LY 25	Bartonella Elizabethae / Rochalimaea elizabethae	Endocarditis. Particularly among homeless IV drug users.
LY 26	Bartonella Grahamii	Endocarditis and Neuroretinitis
LY 27	Bartonella Henselae / Rochalimaea Henselae	Can cause bacteremia, endocarditis, bacillary angiomatosis, and peliosis hepatis. Causes cat-scratch disease.
LY 28	Bartonella Koehlerae	Human infection may be from infected cats.
LY 29	Bartonella Melophagi	Discovered in 2007 and known to infect humans.
LY 30	Bartonella Muris	
LY 31	Bartonella Peromyscus	
LY 32	Bartonella Quintana / Rochalimaea quintana / Rickettsia quintana	Causes trench fever. Can start out as an acute onset of a febrile episode, relapsing febrile episodes, or as a persistent typhoidal illness. Commonly seen are maculopapular rashes, conjunctivitis, headache and myalgias, with splenomegaly being less common. Most patients present with pain in the lower legs (shins), sore muscles of the legs and back, and hyperaesthesia of the shins.
LY 33	Bartonella Rochalimae	Carrion's disease-like symptoms.
LY 34	Bartonella Schoenbuchii	
LY 35	Bartonella Talpae	
LY 36	Bartonella Taylorii	

© Jane Thurnell-Read 1992-2012

Life-Work Potential, Sea View House, Long Rock, Penzance, TR20 8JF, UK
Tel UK: 0844 412 4487; Tel from outside UK: +44 1726 819179
office@lifeworkpotential.com | www.lifeworkpotential.com/

Product Code	Name	Comments
LY 37	Bartonella Tribocorum	
LY 38	Bartonella Vinsonii / Rochalimaea vinsonii	On increase. Causes endocarditis, arthralgia, myalgia, headaches and fatigue.
LY 39	Bartonella Washoensis	May cause fever and myocarditis.
LY 40	Borrelia Afzelii	Has been found in Europe, USA, Singapore, Australia and New Zealand.
LY 41	Borrelia Berbera	Found in Algeria, Tunisia and Libya.
LY 42	Borrelia Burgdorferi	Found in USA, Europe, Australia, New Zealand
LY 43	Borrelia Carteri	Uncommon but has been found in humans in India.
LY 44	Borrelia Caucasica	Found in Europe and Asia.
LY 45	Borrelia Duttonii	Found in Europe and Africa. Causes Central African relapsing fever.
LY 46	Borrelia Garinii	Has been found in Europe.
LY 47	Borrelia Hermsii	Associated with relapsing fever. The primary cause of tick-borne relapsing fever in North America. Also found in Europe.
LY 48	Borrelia Hispanica	Found in Spain, Portugal, Morocco and central Africa.
LY 49	Borrelia Kochis	
LY 50	Borrelia Miyamotoi	Symptoms of relapsing fever. Found in Russia, Japan, Europe and North America.
LY 51	Borrelia Morganii	
LY 52	Borrelia Novyi	Found in the Americas.
LY 53	Borrelia Parkeri	Human infection.
LY 54	Borrelia Persica	Found in Europe and Asia.
LY 55	Borrelia Recurrentis	Found in England, Ireland, USA, Canada, Mexico, Central and South America, central Asia, Africa, and around the Mediterranean.
LY 56	Borrelia Tillaie	Found in Europe.
LY 57	Borrelia Turicatae	Found in Europe.
LY 58	Borrelia Valaisiana	Causes Lyme's disease.
LY 59	Borrelia Venezuelensis	Causes relapsing fever in central and south America.
LY 60	Borrelia Vincentii	Exists normally in the human mouth in low concentrations and safe proportions. Causes severe ulcerating gingivitis (trench mouth); typically found in those with poor oral hygiene but can also occur as a result of stress, cigarette smoking and poor nutrition; also can be found in those with serious illnesses.
LY 61	Ehrlichia Chaffeensis / Human Monocytic Ehrlichiosis	Causative agent of human monocytic ehrlichiosis.

[Buy This Kit](#)